6.0 Reference :

(1) Abu-EI- Ghani.(2000). Vegetation composition of Egyptian Inland saltmarshes, Bot. Bull.Acad.Sin. 41:305-314.

(2) Agoramoorthy, G.Chen. F. A. and Hsu, M.J.(2008) Threat of heavy metals pollution in halophytic and mangrove plant of Tamilnadu, India. Enviro. Pollu. 155,320-326.

(3)Ashraful,M.,Assim, Z. and Ismail's. (2009) Monitoring and assessment of heavy metals levels in littoral sediments from the north eastern part of bay of Bengal coast, Bangladesh. Journal of Industrial pollution control, 25:105-111.

(4) Balasubramanium, T (1991), Coastal Zone biodiversity. UNU International training workshop on methodological for assessing biodiversity in estuaries, mangrove & coastal water.

(5) Breckle, S. W. (2009). Is sustainable agriculture with seawater irrigation realistic ? In: salinity and water stress: improving crop efficiency. Springer, pp. 187-196.

(6) Brink (1993). The coastal ocean progresses effort. Oceanus, 36, 47-49.

(7)Ernst- Detief Schulze –Erwin Beak. Klaus Muller- Hohenstein (2002), Plant Ecology. Ch. 1.7 Pages no. 175-182.

(8)Geddada Mohan Narasimha Rao & Pragada prayaga Murty (2013) Morphological Features of salicornia branchiate Roxb. , Journal of Biological and chemical Research. Vol.30(2) Pages no. 887-891.

(9) GRIGORE Marius-NIcusor, (2012). Halophytes between the fall of civilizations and Biosaline agriculture. EcologicalDisturbances over time, Studiisicomunicari. Stiintelew Naturii. Tom.26. Pages no. 199-204.

(10) Hamdoon A. Mahammed et. Al,(2013), Antioxidant and Quantitative Estimation of Phenolic and Flavonoids of three Halophytic plants Growing in Libya, Journal of pharmacognosy and phytochemistry,Vol 2(3). 89-94.

(11) Hans-werner Koryo et al (2006) strategies of a potential cash crop halophyte (*Beta vulgaris* sp maritima to avoid salt injury, International society for tropical Ecology, Vol 47(2) : 191-200.

(12) ICUN & UNEP, (2003). World database on protected areas. Cambridge, UK.

(13) IP, C.C.M., Li, X. D., Zhang. G. Wai, O. W. H. and Li,Y.S. (2006), Trace metal distribution in sediments of the pearl River Estuary and the surrounding coastal area, south china J. Environ. Pollute. , 147:311-323.

(14) Khot, S.S.(2003). Ecophysiological studies on some halophytes occurring in and aroud marine National park of India. Ph.D. Thesis. Bhavnagar.

(15) Kremer, P. and Van Andel, J. (1995). Evolutionary aspect of life forms in Angiosperms families. Acta Bot.Neerl. 44,469-479.

(16) Mrunalini N. Desai and N. S. Chavan (2011) salt tolerance in seedlings of mangroves Cynometra iripa Kostel. International Journal of applied Biology and pharmaceutical technology. Vol 2(1) : pages no. 219-225.

(17) Nayak, S. Chauhan, H.B.Bahuguna, A & Nath, N. (1996), IRS. IC application for coastal Zone management current science, 70,7

(18) Navjot S. Sodhi and Paul R. Enrich. (2010) conservation Biology for all. OXFORD Univ. press. Ch-2, 27-42.

(19) O'Leary J.W. & Glenn (1994) Global distribution and potential for halophytes. In : Halophytes as a resource for live stock and for rehabilitation of degraded lands. Tasks for vegetation science, 32, pp 7-15.

(20) Rameshkumar S. &Eswaran K (2013) Ecology, utilization and coastal, Management of salt tolerant plants (Halophytes and Mangroves) of MYPAD

coastal regions, Andhrapradesh, India, International journal of Environmental Biology Vol. 3(1) pages no. 1-10.

(21) Robert R. sokal and F. Enrich Rohlf (2009) Introduction to Biostatics, Ch. 1 to 3. Pages no. 46-85.

(22) Roohi Aslam et al. (2011) A Critical reviews on halophytes: salt tolerant Plants. Journal of medicinal plants Research Vol, 5(33) pages no. 7108-7118.

(23) Ray, G.C. (1991), coastal –zone biodiversity patterns. Bioscience, 41, 490-498.

(24) Samiullah and AsghariBanu (2015) Physiological and biochemical analysis of selected halophytes of District Mardan, Pakistan Vol.1(4): 239-243.

(25)S. BHATT, D.G.SHAH& N. DESAI (2009)the mangrove diversity of Purna Estuary, for tropical Ecology. Vol. (50)2. Pages no.- 287-293.

(26)Shigeyuki Baba, Hung Tuck Chan & Sanit Aksornkoae (2013). Useful Products from mangrove & other coastal plants. ISME Mangrove Educational Book series No-3. Okinawa, Japan. ITTO Yokohama, Japan Ch. 5, Pages no. 77-82.

(27)S.sandilyan (2014) Impacts of climate change on Indian Mangroves. Global Journal of Environmental Research, vol. 8(1), pages no.1-10.

(28) Stachowicz, J.J. and Tilman, D. (2005) What species invasions tell us about the relationship between community. Saturation ,species diversity &ecosystem functioning. Pages 41-64. In : D. Sax, J. Stachowicz & s. Gaines (edts.) species invasions: insights in to ecology, evolution& Biogeography. Sinauer, Sunderland, MA.

(29)Shukla, K.H. (2007). Ecophysiological studies on salt tolerance in some halophytes. Ph. D. Thesis, Bhavnagar univ., Bhavnagar.

(30) Tielong Cheng, Jinhi Chen and Jisenshi, Physiological and proteomic analyses of leaves from the halophytes Tangut Nitraria reveals diverse,vol 1(3): 1-298.

(31) Vyas,S. J. (2007) biodiversity of coastal flora in Bhal region part-1. Ph. D. Thesis, Bhavnagar univ. Bhavnagar.

(32)V. Selvam, (2007) Trees and shrubs of the Maldives, FAO Regional office for Asia & the pacific Bangkok, Thailand ch. 4, page no. 177-180.

(33)Weis,p.,Windham,L.,Burke, D.J. and Weis,J.S.(2002).Release into the environment of metals by two vascular salt marsh plant. Mar Environ. Res., 54, 325-329.

(34)Willams, T.P.,Bubb, J. M. and Lester, J. N. (1994a). Metal accumulation within salt marsh environment: a review .Mar. Pollut. Bull.,28(5), 277-290.

(35)Willams, T.P.,Bubb, J. M. and Lester, J. N. (1994b). The occurance and distribution of trae eta n halophytes. Chemosphere,28(6), 1189-1199.

(36) Windham, L., Weis, J.S. and Weis, P. (2003). Uptake and distribution of metals in two dominant salt marsh macrophytes, *spartina alterniflora* (cordgrass) and *Phragmites australis* (common reed) Estuar. Coast shelf sci, 56, 63-72.